

ANORECTAL LYMPHOGRANULOMA VENEREUM: THE FIRST TWO CONFIRMED CASES IN PORTUGAL

R Castro (ritacastro@ihmt.unl.pt)¹, T Baptista², A Vale¹, H Nunes¹, E Prieto¹, K Mansinho², F da Luz Martins Pereira¹

1. Unidade de Doenças Sexualmente Transmitidas (Sexually Transmitted Diseases Unit), Instituto de Higiene e Medicina Tropical (Institute of Hygiene and Tropical Medicine), Universidade Nova de Lisboa (New University of Lisbon), Lisbon, Portugal

2. Serviço de Infecçioologia e Medicina Tropical do Hospital de Egas Moniz (Infectiology and Tropical Medicine Department, Egas Moniz Hospital), Centro Hospitalar de Lisboa Ocidental, Lisbon, Portugal

We describe two cases of lymphogranuloma venereum (LGV) in men who have sex with men in Portugal in 2008. These first two confirmed cases of LGV L2b proctitis in Portugal highlight the need for an enhanced surveillance programme in Portugal.

Lymphogranuloma venereum (LGV) is a sexually transmitted infection (STI) caused by *Chlamydia trachomatis* serovars L1 to L3, which are endemic in tropical regions. However, since 2003, an outbreak of anorectal LGV has emerged in Western Europe [1-4] and in non-Europeans countries [5-7] in men who have sex with men (MSM).

In April 2008, we analysed rectal exudates, urine and blood specimens from two MSM, both infected with human immunodeficiency virus (HIV), that were suspected of having LGV. Both patients presented with an anorectal syndrome. They also had tenesmus and ulcerative lesions, but one of them also suffered from constipation and fever, while the other had inguinal lymphadenopathy and a past infection with *Treponema pallidum*. They were not co-infected with other STIs than LGV and HIV.

One patient was of Portuguese nationality and the other a Brazilian citizen. Both were residents in Portugal and acquired the infection in this country, since they had not travelled abroad six months prior to infection.

IgG antibodies against *C. trachomatis* were high (1:10,000) in both patients, as determined by indirect immunofluorescence (Euroimmun AG, Lübeck,) and *C. trachomatis* serovar L, was identified by real-time polymerase chain reaction (PCR) [8] in the two rectal exudates. Urine samples were negative. The presence of serovar L2b was confirmed by sequencing after amplification of the *omp1* gene by a nested PCR technique [9].

In accordance with the definition of LGV cases [1], the referred patients were confirmed to have LGV infection with the L2b serovar of *C. trachomatis*.

LGV outbreaks have been reported in many countries in Europe, the United States, Canada and Australia [1-7]. In some of these countries, successful enhanced surveillance programmes have been launched, for instance in France [4], the Netherlands [10], and the United Kingdom [3]. Portugal has no surveillance programme; most laboratories do not test for LGV, and not all rectal samples are genotyped.

To our knowledge, these are the first two cases of LGV L2b proctitis described in Portugal. Other cases of LGV have been described in Portugal [11], but those patients had no LGV symptoms and *C. trachomatis* serovar L was only identified in urogenital samples, in contrast to the LGV rectal presentations presently emerging in western Europe.

We therefore feel that there is an urgent need for an enhanced surveillance programme to be launched in Portugal, with a view to alerting clinicians to include LGV in the differential diagnosis of proctitis – especially in HIV-infected MSM. It is equally necessary to inform high risk populations of the signs and symptoms of LGV and of its risks of transmission in view of preventing dissemination of the disease.

Acknowledgements

We thank our colleague Professor Doutor Joao Piedade for helping us with the final identification of *C. trachomatis*, serovar L2b.

References

1. Götz H, Nieuwenhuis R, Ossewaarde T, Thio B, van der Meijden W, Dees J. Preliminary report of an outbreak of lymphogranuloma venereum in homosexual men in the Netherlands, with implications for other countries in western Europe. *Euro Surveill.* 2004;8(4):pii=2367. Available from: <http://www.eurosurveillance.org/ViewArticle.aspx?ArticleId=2367>
2. Liassine N, Caulfield A, Ory G, Restellini A, de Barbeyrac B, Sitavanc R, et al.. First confirmed case of lymphogranuloma venereum (LGV) in Switzerland. *Euro Surveill.* 2005;10(28):pii=2748. Available from: <http://www.eurosurveillance.org/ViewArticle.aspx?ArticleId=2748>
3. Ward H, Martin I, Macdonald N, Alexander S, Simms I, Fenton K et al. Lymphogranuloma venereum in the United Kingdom. *Clin Infect Dis.* 2007;44:26-32.
4. Herida M, de Barbeyrac B, Sednaoui P, Scieux C, Lemarchand N, Kreplak G, et al. Rectal lymphogranuloma venereum surveillance in France 2004-2005. *Euro Surveill.* 2006;11(9):pii=647. Available from: <http://www.eurosurveillance.org/ViewArticle.aspx?ArticleId=647>
5. Kropp RY, Wong T, Canadian LGV Working Group. Emergence of lymphogranuloma venereum in Canada. *CMAJ.* 2005;172(13):1674-6.
6. Tinmouth J, Rachlis A, Wesson T, Hsieh E. Lymphogranuloma venereum in North America: case reports and an update for gastroenterologists. *Clin Gastroenterol Hepatol.* 2006;4(4):469-73.
7. Stark D, van Hal S, Hillman R, Harkness J, Marriott D. Lymphogranuloma venereum in Australia: anorectal *Chlamydia trachomatis* serovar L2b in men who have sex with men. *J Clin Microbiol.* 2007;45(3):1029-31.

8. Kivi M, Koedijk FD, van der Sande M, van de Laar MJ. Evaluation prompting transition from enhanced to routine surveillance of lymphogranuloma venereum (LGV) in the Netherlands. *Euro Surveill.*; 2008.13(14);pii=8087. Available from: <http://www.eurosurveillance.org/ViewArticle.aspx?ArticleId=8087>
9. Spaargaren J, Fennema HS, Morr  SA, de Vries HJ, Coutinho RA. New lymphogranuloma venereum Chlamydia trachomatis variant, Amsterdam. *Emerg Infect Dis.* 2005;11(7):1090-2.
10. Yang CL, Maclean I, Brunham RC. DNA sequence polymorphism of the Chlamydia trachomatis omp1 gene. *J Infect Dis.* 1993;168(5):1225-30.
11. ESSTI alert. LGV Cases Reported in Denmark and Portugal. London: Health Protection Agency, European Surveillance of Sexually Transmitted Infections; 2007 Jun. Available from: http://209.85.129.132/search?q=cache:bjs1o279wHgJ:www.essti.org/docs/ESSTI_Newsletter_issue_5.pdf+ESSTI+alert+portugal&hl=pt-PT&ct=clnk&cd=2&gl=pt

This article was published on 11 December 2008.

Citation style for this article: Castro R, Baptista T, Vale A, Nunes H, Prieto E, Mansinho K, da Luz Martins Pereira F. Anorectal lymphogranuloma venereum: the first two confirmed cases in Portugal. *Euro Surveill.* 2008;13(50);pii=19060. Available online: <http://www.eurosurveillance.org/ViewArticle.aspx?ArticleId=19060>